



NETWORK-RED

734H000122

JIMTOWN NETWORK-RED-ET

MARK *RC X SALVATORE RC X ANDIRON *RC

H0840M003150919056 | DOB 10/29/2019

Kappa-Casein AB | Beta-Casein A2A2

Haplotypes HH1F HH2F HH3F HH4F HH5F HCDF HH6F

Genetic Codes TC TD TL TP TV TY

aAa 234 | DMS 234,123 | EFI 10.5% | RHA 99%

Current Proof USA-202112

CDCB & HA-USA Genetic Evaluations 12/2021

CONFORMATION Based on 0 Daughters in 0 Herds (78% Rel)

| | PTAT | 1.87 | Udder Composite | 1.45 | | | | |
|----------------|-----------------------|------|----------------------|------|---|---|-------|-----------|
| | Body Weight Composite | 0.90 | Foot & Leg Composite | 0.50 | | | | |
| | | -2 | -1 | 0 | 1 | 2 | | |
| Stature | | | | | | | +1.61 | Tall |
| Strength | | | | | | | +1.02 | Strong |
| Body Depth | | | | | | | +1.05 | Deep |
| Dairy form | | | | | | | +1.01 | Open |
| Rump Angle | | | | | | | -0.46 | High Pins |
| Thurl Width | | | | | | | +1.12 | Wide |
| R. Legs-S View | | | | | | | +0.85 | Sickle |
| R. Legs-R View | | | | | | | +0.86 | Straight |
| Foot Angle | | | | | | | +0.95 | Steep |
| F&L Score | | | | | | | +0.80 | High |
| F. Udder Att. | | | | | | | +2.04 | Strong |
| R. Udder Ht. | | | | | | | +1.91 | High |
| R. Udder Wid. | | | | | | | +2.31 | Wide |
| Udder Cleft | | | | | | | +0.90 | Strong |
| Udder Depth | | | | | | | +1.22 | Shallow |
| F. Teat Place | | | | | | | +0.90 | Close |
| R. Teat Place | | | | | | | +1.03 | Close |
| Teat Length | | | | | | | +0.61 | Long |

PEDIGREE

| | |
|------|----------------------------|
| SIRE | WILDER MARK |
| DAM | JIMTOWN SAL NAYELI-RED-ET |
| MGS | MR SALVATORE RC-ET |
| MGD | JIMTOWN AND NATALYA-RED-ET |
| MGGS | MR ADERYN ANDIRON-ET |
| MGGD | JIMTOWN SHAN NATINE-ET |

| | | | |
|-----|------|------|-----|
| TPI | 2786 | NM\$ | 737 |
|-----|------|------|-----|

PRODUCTION

| | | | | |
|---------|-----------|---------|------|-------|
| Milk | +1525 Lbs | 80% Rel | CM\$ | \$740 |
| Protein | +46 Lbs | 0.00% | FM\$ | \$731 |
| Fat | +75 Lbs | +0.06% | GM\$ | \$682 |

Based on 0 Daughters in 0 Herds (100% US Daughters)

| |
|--------------------------------|
| Milk Production - US daughters |
| Fat Yield - US daughters |
| Protein Yield - US daughters |

HEALTH TRAITS

| | | | |
|---------------------|-------|-------------------------|-------|
| Productive Life | +4.6 | Daughter Pregnancy Rate | +0.1 |
| Cow Conception Rate | 0.1 | Heifer Conception Rate | 0.3 |
| Cow Livability | +0.5 | Heifer Livability | +0.8 |
| Somatic Cell Score | +2.86 | Mastitis | 0.3% |
| Metritis | 1.9% | Ketosis | 2.0% |
| Displaced Abomasum | 0.7% | Retained Placenta | 0.4% |
| Feed Saved | -121 | Milk Fever | +0.1% |
| Zoetis DWP\$ | \$609 | Zoetis WT\$ | \$-27 |
| Zoetis CW\$ | \$-17 | | |

CALVING TRAITS

| | | | | | |
|-------------------|------|---------|---------------------|------|---------|
| Sire Calving Ease | 1.9% | 71% Rel | Sire stillbirth | 5.7% | 63% Rel |
| Dtr Calving Ease | 2.1% | 71% Rel | Daughter stillbirth | 4.6% | 65% Rel |